

## Original Article

*Iran J Ped Hematol Oncol. 2016, Vol6.No3,172-181***Gene Expression and Promoter Methylation Status of VHL, Runx-3, E-cadherin, P15 and P16 Genes During EPO-Mediated Erythroid Differentiation of CD34+ Hematopoietic Stem Cells**

Fatemeh Eskandari MSc<sup>1</sup>, Hoda Pourkarim MSc<sup>2</sup>, Mehdi Sahmani PhD<sup>3</sup>, Mehdi Goudarzi PhD<sup>4</sup>, Naser Mobarra PhD<sup>5</sup>, Ali Dehghanifard PhD<sup>6</sup>, Gholamreza Khamisipour PhD<sup>7</sup>, Nasim Kalantari MSc<sup>1</sup>, Hanieh Rahmani MSc<sup>8</sup>, Mehdi Azad PhD<sup>9,\*</sup>

1. Department of Hematology, School of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Department of Hematology, Allied Medical School, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Clinical Biochemistry, Cellular and Molecular Research Center, Qazvin, Iran

4. Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Science, Tehran, Iran

5. Stem cell Research Center, School of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

6. Sarem Cell Research Center, Sarem Women's Hospital, Tehran, Iran

7. Department of Hematology, Faculty of Allied Medicine, Bushehr University of Medical Sciences, Bushehr, Iran

8. Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

9. Department of Medical laboratory sciences, Faculty of Allied Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

\*Corresponding Author: Mehdi Azad. PhD. Department of Medical laboratory sciences, Faculty of Allied Medicine (Hematology), Qazvin University of Medical Sciences, Qazvin, Iran. Email Address: haematologicca@gmail.com.

**Received:** 28 November 2015

**Accepted:** 12 March 2016

**Abstract**

**Background:** VHL (von Hippel-Lindau), Runx-3 (Runt-related transcription factor 3), E-cadherin (Epithelial cadherin), P15 (INK4a, cyclin dependent kinase inhibitor), and P16 (INK4b) genes are essential in hematopoiesis. The aim of this study was to explore the correlation between gene expression and promoter methylation in CD34+ stem cells before and after differentiation to erythroid lineage.

**Materials and Methods:** CD34+ hematopoietic stem cells were separated from umbilical cord blood using MidiMacs (positive selection) system. Expanded CD34+ stem cells were differentiated into erythroid lineage with human recombinant erythropoietin (EPO). DNA extraction was done by QIAamp DNA Mini Kit. RNA was extracted using RNase Mini plus Kit. MSP (Methylation specific PCR) technique was done for methylation assay. Methylation status and expression assay was done for VHL, Runx-3, E-cadherin, P15, and P16 genes on both CD34+ stem cells and differentiated erythroid cells.

**Results:** The results showed that, before differentiation, P15 had comparative methylation pattern and average expression and it remained unchanged after differentiation ( $p=0.01$ ). concerning P16, results revealed no methylation pattern and complete expression in absence of EPO and with EPO it changed to comparative status ( $p=0.01$ ). E-cad and Runx-3 genes had relative methylation pattern and fully expression before and after differentiation but their expression after that, was increased and decreased Respectively ( $p=0.04$ ). VHL gene had no significant methylation status before or after differentiation and its expression was complete ( $p=0.01$ ).

**Conclusion:** The obtained results indicated that promoter methylation of P15, P16, VHL, Runx3 and E-cad was one of the definitive expression control mechanism of these genes.

**Key words:** Differentiation, EPO, Hematopoietic Stem Cells

**Introduction**

Studies about cell differentiation have led to access general mechanisms of regulating and specific factors of gene expression. In fact intracellular signaling and gene expression regulators are the most impressive factor for primary

differentiation process in the cell (1). So the requirement for specific elements like cell cycle controlling molecules or cytokines is quite understandable. In mammalian cells DNA is methylated only at cytosine located 5' to guanosine in CpG